

First record of *Camarops petersii* from eastern Europe

EUGENE S. POPOV^{*A} & SERGEY V. VOLOBUEV^{*B}

Komarov Botanical Institute of the Russian Academy of Sciences, Laboratory of Systematics and
Geography of Fungi, Saint Petersburg 197376, Russia

* CORRESPONDENCE TO: ^A epopov@binran.ru ^B sergvolobuev@binran.ru

ABSTRACT—The stromatic ascomycete *Camarops petersii* was recorded for the first time in Eastern Europe. It was found growing on *Quercus robur* wood, on a fallen trunk of a lightning stuck tree in Bryansk Oblast, next to the western border of Russia. The morphological description, data on the distribution and habitat are provided. The modern distribution of *C. petersii* is confined mostly to the refugia of tertiary relict floras. All known European populations of *Camarops petersii* (including our own) are from north of the Alps, possibly established relatively recently due to post-glacial colonization of the broadleaf forests. Future *C. petersii* populations may be found in the Mediterranean area and the Caucasus.

KEY WORDS—*Boliniaceae*, *Boliniales*, fungal ecology, mycogeography, *Sordariomycetes*

Introduction

Camarops P. Karst. (*Boliniaceae*, *Boliniales*) is a genus of stromatic *Sordariomycetes* characterized by massive, erumpent, usually dark colored and carbonized stromata, evanescent paraphyses and asci, and relatively small, brown-colored, non-septate ascospores with a germ pore (Nannfeldt 1972). The genus includes 25 species according to Index Fungorum (2019).

Camarops petersii is well distinguished from all other species of the genus because the immature stroma is covered by a thick, membranaceous veil-like structure that breaks down later except at the margin. The species was originally described based on two collections from Alabama, (USA) and Cuba (Berkeley 1868) and for a long time was believed to occur only in the western hemisphere until it was discovered in Korea in 1930 (Iwade 1944) and later in Germany (Nuss & Hilber 1977).

In the present paper, we discuss the first record of *Camarops petersii* in eastern Europe based on a collection made by the second author during a survey of wood-inhabiting fungi in Bryansk Oblast (Russia).

Material & methods

The field survey was organized by the second author in August 2018 during a multi-year survey of the wood-inhabiting fungi of the Non-Chernozem Zone of Russia (Popov 2013, Volobuev 2013, Popov & Volobuev 2014). Voucher specimens were deposited in the Mycological Herbarium of the Komarov Botanical Institute of the Russian Academy of Sciences, Saint Petersburg, Russia (LE).

Global distribution data for *Camarops petersii* was taken from literature and the Global Biodiversity Information Facility portal (GBIF 2019), which cited 112 occurrences on 7 April 2019.

The description of macroscopic characters was based on fresh and dry material and photographs in situ. Microscopic features were observed and measured from dried material squash mounted in tap water for light microscopy (Zeiss Axio Imager A1 microscope). Melzer's reagent (MLZ) was used to check the amyloid reaction of hymenial structures. Dimensions of spores were derived from 20 measures. The images of microstructures were captured with an AxioCam MRC 5 digital camera on the microscope equipped with bright-field and differential interference contrast (DIC) optics. Images were edited by using AxioVision SE64 ver. 4.9.1 software.

DNA extraction, amplification, and sequencing followed Zmitrovich & al. (2019). The fungal specific primers ITS1F and ITS4 were used to sequence the ITS rDNA gene region (White & al. 1990, Gardes & Bruns 1993).

Taxonomy

Camarops petersii (Berk. & M.A. Curtis) Nannf.,

Svensk Bot. Tidskr. 66(4): 366 (1972).

PL. 1

STROMATA superficial on wood, individual, 3–5 cm in diam, ≤1 cm high, flattened-turbinate to discoid, centrally attached, woody, dull brown in cross section, initially wrapped in a whitish, brown flecked veil, which irregularly cracks and breaks down when mature to expose a black and shiny stromatal surface; PERITHECIA immersed in the upper part of stromata at different depth (polystichous), obclavate, 1–1.5 × 0.4–0.5 mm, with long necks ending in papillate ostioles.

ASCI 52–65 × 4.8–6.3 µm, cylindrical, stipitate, with an inamyloid apical ring, 8-spored, evanescent and often absent in fully mature perithecia; ASCOSPORES 6.0–8.4 × 3.3–4.5 µm, uniseriate, 1-celled, ellipsoid to obovate, slightly pointed at one end, olive brown, with two large lipid drops and a germ pore at the



PLATE 1. *Camarops petersii* (LE 323478). A. Fresh mature stroma *in situ* showing disrupted veil and exposed stromatal surface; B. Stroma in cross section with polystichous perithecia in upper part; C. Ascospores (position of germ pore marked with an asterisk). Scale bars: A, B = 1 cm; C = 5 μ m.

tapered end; PARAPHYSES cylindrical, 1.5–2 μ m diam., hyaline, septate, longer than asci and evanescent.

MATERIAL EXAMINED: RUSSIA, BRYANSK OBLAST: Suzemskiy district, Nerusso-Sevnyi protected forest area, 'Rybnitsa' natural boundary, 52°24'32"N 34°05'00"E, floodplain, oak-dominated deciduous forest with *Quercus robur*, *Acer platanoides*, *Fraxinus excelsior*, on decorticated trunk of *Q. robur* lying on the ground, 30 August 2018, coll. S. Volobuev (LE 323478; GenBank MN564897).

Discussion

Camarops petersii grows on fallen and usually decorticated logs of *Quercus* spp, more rarely of *Castanea*, *Castanopsis*, and *Zelkova* (Nannfeldt 1972, Nuss & Hilber 1977, Doi & Nunomura 1980). Our collection was also found on wood of a barkless, recently fallen trunk that was about 60 cm in diam, and comparatively hard (a knife could not penetrate more than 2–3 cm in depth).

Camarops petersii appears to be widely distributed in North America, where it has been frequently reported from broadleaf forests of the Appalachians and Central Lowlands west to Kansas and as far north as Vermont and Wisconsin (Ellis & Everhart 1888, Shear 1923, Heagle & French 1972, Nannfeldt 1972, Fergus 1973, Horn 1984, Vasilyeva & al. 2007). Moreover, 77 of 112 occurrences of this species in the GBIF portal are from North America (GBIF 2019).

Infrequent findings have been reported from temperate broadleaf and mixed forests in Asia: South Korea (Iwade 1944), Russian Far East (Nannfeldt 1972), Japan (Doi & Nunomura 1980), Taiwan (GBIF 2019), northern Iran (Arefipoor & al. 2018), and northwestern Himalayas (Dargan 1976; Dargan & Thind 1984); and in Europe, from Germany, Austria, and France (Nuss & Hilber 1977, Urban 2016, MycoDB 2019). The modern disjunct distribution of *C. petersii* is thus confined mostly to the refugia of tertiary relict floras as defined by Milne & Abbott (2002).

In Europe, during most of the last glacial period, the temperate forest species, including summer-green species of the *Quercus*, were restricted to small refugia. Those were situated mostly south of 45°N along the Mediterranean coast and around the Black and Caspian Seas (Leroy & Arpe 2007, Svenning & al. 2008, Tzedakis & al. 2013). At that time, most of the area north of the Alps was covered with tundra and cold steppes (Gerasimov & Velichko 1982, Binney & al. 2017). The spread of deciduous *Quercus* species from these refugia to the Central European mountains took place in the late-glacial interstadial (13–11 thousand years ago), and oaks spread into Northern and Eastern Europe only in the Holocene (Brewer & al. 2002, Birks & Tinner 2016). It is likely that European populations of *Camarops petersii* were recently established as a result of post-glacial colonization of the broadleaf forests. We suppose that *C. petersii* still could be found in the Mediterranean area and the Caucasus. But this supposition needs to be verified by mycologists in those regions.

A BLAST search revealed that the ITS sequence newly generated from our material (GenBank MN564897) is identical with those of the North American collection of *C. petersii* MICH 340497 (GenBank MT913624).

Acknowledgments

The authors are grateful to Dr. Elena Sitnikova (deputy director of the Bryanskiy Les Nature Reserve) for great assistance with field work management. This study was carried out within the framework of the institutional research project Biodiversity, ecology, structural and functional features of fungi and fungus-like protists (AAAA-A19-119020890079-6) of the Komarov Botanical Institute RAS using equipment of its Core Facility Center [Cell and Molecular Technologies in Plant Science]. The authors also thank Drs. Viktor Kučera (Slovak Academy of Sciences, Bratislava, Slovakia) and Luis Quijada (Harvard University Herbaria, Cambridge MA, USA) for presubmission review.

Literature cited

Arefipoor MR, Farashiani E, Mousazadeh SA, Zamani SM, Aghajani H, Rafeijahed R. 2018. Introduction of some fungal species saproxylic in the forest of Golestan province. 77, in:

- Proceedings of the 1st International Congress and the 2nd National Conference on Biotechnology of Medicinal Plants and Mushrooms, University of Zanjan, August 27–28. Zanjan, University of Zanjan [Accessed 03 April 2019]: [http://conf.isc.gov.ir/FileForDownload/files/IBMPM97/Abstract English .pdf](http://conf.isc.gov.ir/FileForDownload/files/IBMPM97/Abstract%20English.pdf).
- Berkeley MJ. 1868 ["1869"]. On a collection of fungi from Cuba. Part. II., including those belonging to the families *Gasteromycetes*, *Coniomycetes*, *Hyphomycetes*, *Phycomycetes*, and *Ascomycetes*. Journal of the Linnean Society, Botany 10(46): 341–392.
- Binney H, Edwards M, Macias-Fauria M, Lozhkin A, Anderson P, Kaplan JO, Andreev A, Bezrukova E, Blyakharchuk T, Jankovska V. 2017. Vegetation of Eurasia from the last glacial maximum to present: Key biogeographic patterns. Quaternary Science Reviews 157: 80–97. <https://doi.org/10.1016/j.quascirev.2016.11.022>
- Birks HJB, Tinner W. 2016. Past forests of Europe. 36–39, in: J San-Miguel-Ayaz J & al. (eds). European Atlas of Forest Tree Species. Luxembourg, Publication Office of the European Union.
- Brewer S, Cheddadi R, De Beaulieu JL, Reille M. 2002. The spread of deciduous *Quercus* throughout Europe since the last glacial period. Forest Ecology and Management 156(1–3): 27–48. [https://doi.org/10.1016/S0378-1127\(01\)00646-6](https://doi.org/10.1016/S0378-1127(01)00646-6)
- Dargan JS. 1976. Studies on the *Xylariaceae* of North Western Himalayas. PhD dissertation, Department of Botany, Panjab University, Chandigarh.
- Dargan JS, Thind KS. 1984. Genus *Peridoxylon* Shear in India. Journal of Indian Botanical Society 63(2): 177–180.
- Doi Y, Nunomura K. 1980. *Camarops petersii* (Berk. & Curt.) Nannf. found in Japan. Bulletin of the Natural Science Museum, ser. B, 6: 97–100.
- Ellis JB, Everhart BM. 1888. Synopsis of the North American species of *Hypoxylon* and *Nummularia*. Journal of Mycology 4(4–5): 38–44.
- Fergus CL. 1973. *Peridoxylon petersii* on oak in Pennsylvania. Mycologia 65(5): 1199–1200. <https://doi.org/10.2307/3758297>
- GBIF. 2019. *Camarops petersii* (Berk. & M.A. Curtis) Nannf. GBIF Occurrence Download. <https://doi.org/10.15468/dl.ufa501> (accessed 07 April 2019).
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes—application to the identification of mycorrhizae and rusts. Molecular Ecology 2: 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Gerasimov IP, Velichko AA. 1982. Paleogeography of Europe during the last one hundred thousand years. Moscow, Nauka
- Heagle AS, French DW. 1972. *Peridoxylon petersii* on oak in Minnesota. Mycologia 64(6): 1349–1351. <https://doi.org/10.2307/3757975>
- Horn BW. 1984. Fertile stromata of *Camarops petersii* in culture. Mycologia 76(5): 956–959. <https://doi.org/10.2307/3793156>
- Index Fungorum. 2019. www.indexfungorum.org/Names/Names.asp (accessed 07 April 2019).
- Iwade I. 1944. Studies on Japanese fungi. I. Several new and rare species of fungi. Bulletin of the Tokyo University Forests 33: 49–64.
- Leroy SA, Arpe K. 2007. Glacial refugia for summer-green trees in Europe and south-west Asia as proposed by ECHAM3 time-slice atmospheric model simulations. Journal of Biogeography 34(12): 2115–2128. <https://doi.org/10.1111/j.1365-2699.2007.01754.x>
- Milne RI, Abbott RJ. 2002. The origin and evolution of Tertiary relict floras. Advances in Botanical Research 38: 281–314. [https://doi.org/10.1016/S0065-2296\(02\)38033-9](https://doi.org/10.1016/S0065-2296(02)38033-9)
- Mycodb. 2019. Liste des récoltes pour *Camarops petersii*. www.mycodb.fr/recolte.php?source=popup&action=list&genre=Camarops&espece=petersii (accessed 07 April 2019).

- Nannfeldt JA. 1972. *Camarops* Karst. (*Sphaeriales*–*Boliniaceae*) with special regard to its European species. *Svensk Botanisk Tidskrift* 66: 335–376.
- Nuss I, Hilber R. 1977. *Camarops petersii* (Berk. et Curt.) Nannf. Erstnachweis für Europa — und weitere *Camarops*-Arten. *Zeitschrift für Pilzkunde* 43: 217–236.
- Popov ES. 2013. New records of some rare species in the family *Hyaloscyphaceae* (*Ascomycota*, *Helotiales*) from Central Russia. *Novosti Sistematiki Nizshikh Rasteniy* 47: 135–142.
- Popov ES, Volobuev SV. 2014. New data on wood-inhabiting macromycetes in key protected areas of the south-western part of the Non-Chernozem Zone. *Mikologiya i Fitopatologiya* 48(4): 229–237.
- Shear CL. 1923. Life histories and undescribed genera and species of fungi. *Mycologia* 15(3): 120–131. <https://doi.org/10.2307/3753058>
- Svenning JC, Normand S, Kageyama M. 2008. Glacial refugia of temperate trees in Europe: insights from species distribution modelling. *Journal of Ecology* 96(6): 1117–1127. <https://doi.org/10.1111/j.1365-2745.2008.01422.x>
- Tzedakis PC, Emerson BC, Hewitt GM. 2013. Cryptic or mystic? Glacial tree refugia in northern Europe. *Trends in Ecology & Evolution* 28(12): 696–704. <https://doi.org/10.1016/j.tree.2013.09.001>
- Urban A. 2016. First record of the dog's nose fungus, *Camarops petersii*, in the old-growth forest reserve Johannserkogel, Vienna, Austria. *Austrian Journal of Mycology* 25: 225–229.
- Vasilyeva LN, Stephenson SL, Miller AN. 2007. Pyrenomycetes of the Great Smoky Mountains National Park. IV. *Biscogniauxia*, *Camaropella*, *Camarops*, *Camillea*, *Peridoxylon* and *Whalleya*. *Fungal Diversity* 25: 219–231.
- Volobuev S. 2013. The first record of *Jaapia ochroleuca* (*Basidiomycota*) in the European part of Russia. *Novosti Sistematiki Nizshikh Rasteniy* 47: 68–73.
- White TJ, Bruns T, Lee S, Taylor JW. 1990. Amplification of direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis & al. (eds). *PCR protocols: a guide to methods and applications*. San Diego, Academic Press. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Zmitrovich IV, Volobuev SV, Dudka VA, Zhukova EA, Sidelnikova MV, Bondartseva MA. 2019. *Ganoderma applanatum* (*Polyporales*, *Basidiomycota*) at the Saint Petersburg area. *Mikologiya i Fitopatologiya* 53(6): 354–362. <https://doi.org/10.1134/S0026364819060084>